

WHAT IS CLAIMED IS:

1. A method of screening for a transdominant bioactive agent, said method comprising steps:  
expressing a molecular library of randomized nucleic acids as a plurality of isolated  
corresponding randomized translation products in a first plurality of cells, each of said nucleic  
acids comprising a different nucleotide sequence;

5 screening a second plurality of cells for a cell exhibiting a changed physiology in response  
to the presence of a transdominant translation product of said plurality of isolated corresponding  
randomized translation products, wherein said translation product is expressed with a fusion  
partner, synthetic or heterologous to said first plurality of cells, comprising a localizing signal  
sequence capable of constitutively localizing said translation product to a predetermined  
10 subcellular locale, secretory and membrane-anchoring signal sequences capable of localizing said  
translation product to the plasma membrane, or a secretory signal sequence capable of effecting  
the secretion of said translation product.

detecting said cell;

15 isolating at least one of said cell and said transdominant translation product, wherein said  
transdominant translation product is a transdominant bioactive agent.

2. A method according to claim 1, wherein said translation products are presented on the  
extracellular surface of said first plurality of cells.

20 3. A method according to claim 1, wherein said translation products are secreted from said first  
plurality of cells.

4. A method according to claim 1, wherein said first and second plurality of cells are different.

25 5. A method according to claim 1, wherein said expressing step further comprises introducing  
said library into said cells.

6. A method according to claim 1, wherein said expressing step further comprises introducing

said library into said cells using retroviral vectors.

7. A method of screening for a transdominant extracellularly bioactive agent, said method comprising steps:

expressing a molecular library of randomized nucleic acids as a plurality of isolated  
5 corresponding randomized translation products in a first plurality of cells, each of said nucleic acids comprising a different nucleotide sequence;

screening a second different plurality of cells for a cell exhibiting a changed physiology in response to the presence of a transdominant translation product of said plurality of isolated  
10 corresponding randomized translation products, wherein said translation product is expressed with a fusion partner, synthetic or heterologous to said first plurality of cells, comprising a secretory and membrane-anchoring signal sequences capable of localizing said translation product to the extracellular surface of the plasma membrane, or a secretory signal sequence capable of effecting the secretion of said translation product.

detecting said cell;

15 isolating at least one of said cell and said transdominant translation product, wherein said transdominant translation product is a transdominant extracellularly bioactive agent; wherein said expressing step comprises introducing said library into said cells using retroviral vectors.

ADD A17